


## Porphyrin–Polyyne [3]- and [5]Rotaxanes

Daniel R. Kohn, Levon D. Movsisyan, Amber L. Thompson, and Harry L. Anderson\*<sup>1b</sup>

University of Oxford, Department of Chemistry, Chemistry Research Laboratory, Oxford OX1 3TA, United Kingdom

### **S** Supporting Information


**ABSTRACT:** Porphyrin–polyyne [3]- and [5]rotaxanes have been synthesized by condensing aldehyde–rotaxanes with pyrrole or dipyrromethane. The crystal structure of a [3]rotaxane shows that the macrocycles adopt compact conformations, holding the hexaynes near the porphyrin core, and that the phenanthroline units form intermolecular  $\pi$ -stacked dimers in the solid. Fluorescence spectra reveal singlet excited-state energy transfer from the threaded hexayne to the porphyrin, from the phenanthroline to the porphyrin, and from the phenanthroline to the hexayne.



Macrocycles consisting entirely of  $sp$ -hybridized carbon are intriguing synthetic targets.<sup>1–3</sup> Cyclocarbons have been studied in the gas phase,<sup>1,2,4</sup> but they have not yet been isolated or characterized in solution, despite attempts by Diederich, Tobe, Rees, and co-workers.<sup>1–3</sup> Advances in active metal template synthesis<sup>5</sup> provide access to an expanding diversity of interlocked molecules, including threaded polyynes,<sup>6–11</sup> and it has been found that linear chains of  $sp$ -hybridized carbon atoms (both polyynes and cumulenes) can be stabilized by threading them through macrocycles to form rotaxanes.<sup>11–13</sup> As part of a project directed toward the synthesis of cyclocarbon catenanes, we are exploring strategies for positioning several polyynyl rotaxane units around a central molecular hub, so that the polyynes can then be linked together to form a threaded cyclocarbon. Here, we present the first step in this endeavor: the synthesis of a [5]rotaxane in which four hexaynes are positioned around a porphyrin core. While the primary motivation for this work was to explore approaches to cyclocarbon catenanes, this study also provides insights into the interactions between the singlet excited states of polyynyl and porphyrin chromophores.

The initial target of this work was the porphyrin [5]rotaxane **P5Ra** shown in Figure 1. All attempts at the synthesis of this [5]rotaxane failed, presumably as a consequence of the steric bulk of the four [2]rotaxane substituents. However, we successfully synthesized the analogous porphyrin [5]rotaxane **P5Rb** with *p*-phenylene spacers at the *meso*-positions and the corresponding porphyrin [3]rotaxanes **P3Ra** and **P3Rb** (Scheme 1). Here, we present the synthesis of these rotaxanes and an analysis of their fluorescence behavior, together with the crystal structure of a porphyrin [3]rotaxane **P3Ra**. This structure shows that the macrocycles adopt compact conformations due to the predominantly gauche geometries of the  $O-CH_2CH_2-O$  links, which increases the effective steric bulk of the rotaxane substituents and explains why the [5]rotaxane **P5Ra** could not be synthesized.

Our synthetic approach starts with the macrocyclic aldehyde **Ma** (Scheme 1).<sup>14,15</sup> Active metal template Cadiot–Chodkie-




**Figure 1.** Structure of target [5]rotaxane **P5Ra**.

wicz coupling<sup>7,11</sup> of supertrityl triene  $Tr^*C_6H$  and bromotriene  $Tr^*C_6Br$  [where  $Tr^*$  is tris(3,5-di-*tert*-butylphenyl)methyl<sup>16</sup>] in the presence of the copper(I) complex of **Ma** gave the [2]rotaxane aldehyde **2Ra** in 58% yield. A wide variety of conditions were tested for the reaction of **2Ra** with pyrrole in attempts to prepare **P5Ra**, but they all failed to give even a trace of this target [5]rotaxane, whereas control reactions of aldehyde **Ma** produced porphyrin **PM4a** in good yield. In contrast, reaction of the rotaxane aldehyde **2Ra** with dipyrromethane gave the porphyrin [3]rotaxane **P3Ra** in 62% yield. These results

**Received:** November 26, 2016

**Published:** January 5, 2017


Scheme 1. Polyyrotaxane and Porphyrin Synthesis<sup>a</sup>

<sup>a</sup>Porphyrins **PM4a**, **PM2a**, **P3Ra**, **P5Ra**, **PM4b**, **PM2b**, **P3Rb**, and **P5Rb** were isolated as zinc complexes after treatment with  $\text{Zn}(\text{OAc})_2 \cdot 2\text{H}_2\text{O}$  in methanol/chloroform.

imply that formation of **P5Ra** is blocked by the steric bulk of the rotaxane units, so we investigated the analogous compounds derived from macrocycle **Mb**, which has a *p*-phenylene spacer between the aldehyde functionality and the macrocycle. Increasing the distance between the bulky rotaxane unit and the porphyrin hub made it possible to synthesize [5]rotaxane **P5Rb** (10% yield isolated).

The porphyrin [3]- and [5]rotaxanes, and reference porphyrins, were fully characterized as their zinc complexes **Zn-P3Ra**, **Zn-P3Rb**, **Zn-P5Rb**, **Zn-PM2a**, **Zn-PM4a**, **Zn-PM2b**, and **Zn-PM4b** (see the Supporting Information). Crystals of **Zn-P3Ra** were grown by diffusion of methanol vapor into a solution in chloroform and analyzed by single-crystal X-ray diffraction.<sup>17</sup> The crystal structure has two molecules of

Zn–P3Ra in the asymmetric unit; both rotaxane molecules have similar conformations with methanol coordinated to the zinc centers. The phenanthroline units of each rotaxane form intermolecular  $\pi$ -stacked dimers with crystallographically equivalent rotaxanes, forming infinite strands of interactions, as illustrated in Figure 2. The oligoethylene glycol chains of the



**Figure 2.** Projection of the structure of the [3]rotaxane Zn–P3Ra from X-ray crystallography. This figure only shows one of the two molecules in the asymmetric unit.


macrocycles adopt compact conformations and are mostly gauche ( $\text{O}-\text{CH}_2\text{CH}_2-\text{O}$  torsional angle  $40-80^\circ$ ) with the polyene chains held close to the porphyrin units. Thus, all four polyene chains in the asymmetric unit make van der Waals contacts with the  $\beta$ -pyrrole protons of the porphyrins (sp-C $\cdots$ H distance: 2.9–3.1 Å). The hexayne chains have slightly curved geometries, similar to those reported in crystal structures of other polyynes.<sup>9,11,16,18–20</sup>

The family of rotaxanes synthesized during this study provides a unique opportunity to investigate photophysical interactions between the excited states of polyene and porphyrin chromophores. The photophysical behavior of polyynes is particularly interesting because their absorption spectra are dominated by strong transitions to higher excited states at 280–320 nm ( $S_0-S_n$ ,  $\epsilon \approx 3 \times 10^5 \text{ M}^{-1} \text{ cm}^{-1}$ ); absorption into the first excited state is dipole-forbidden and can occasionally be observed as a very weak band at 350–450 nm ( $S_0-S_1$ ,  $\epsilon \approx 500 \text{ M}^{-1} \text{ cm}^{-1}$ ).<sup>16,18</sup> Polyynes also exhibit remarkably fast intersystem crossing ( $S_1-T_1$ ), which usually prevents observation of fluorescence.

The absorption spectra of rotaxanes Zn–P3Ra, Zn–P3Rb, and Zn–P5Rb demonstrate that threading causes very little perturbation to the electronic structure of the component chromophores. Thus, the absorption bands of the polyene (280–320 nm), the phenanthroline macrocycle (250–380 nm), and porphyrin (400–450 nm Soret band and 530–560 nm Q-band) are readily identified in the rotaxanes. Comparison with simple reference porphyrin derivatives (*meso*-tetraphenyl porphyrin for Zn–P5Rb and Zn–PM4a; 5,15-diaryl porphyrin for Zn–P3Ra, Zn–P3Rb, Zn–PM2a, and Zn–PM2b) shows that the polyene and phenanthroline macrocycle units have no effect on the fluorescence quantum yield, measured on excitation of the porphyrin Soret band at around 415 nm. This

demonstrates that there is no significant energy migration from the porphyrin to the polyene or to the macrocycle.

Fluorescence excitation spectra of the rotaxanes, measured by recording emission from the porphyrin Q-band at around 640 nm, show clear features from excitation into the polyene and phenanthroline macrocycle components, as illustrated for Zn–P3Ra and Zn–PM2a in Figure 3, indicating that both of these



**Figure 3.** (a) Absorption spectra (continuous line) and excitation spectra (dashed line) of Zn–P3Ra (red) and Zn–PM2a (blue); the spectral intensities are normalized to 1.0 at the Q-band, and the excitation spectra are recorded for emission at 639 nm (porphyrin Q-band). Solvent:  $\text{CH}_2\text{Cl}_2$ . (b) Jablonski diagram showing the energy-transfer processes.

chromophores transfer energy to the porphyrin. A detailed examination of the absorption spectra, fluorescence spectra, and excitation spectra of the whole family of compounds allowed us to determine the quantum yields of excited-state energy transfer (EET) from the polyene to the porphyrin and from the phenanthroline macrocycle to the porphyrin as summarized in Table 1 (see the SI for full details).

**Table 1. Energy-Transfer Efficiencies  $\phi_{\text{EET}}$**

compd	$\phi_{\text{EET}}(\text{macrocycle} \rightarrow \text{porphyrin})$	$\phi_{\text{EET}}(\text{polyene} \rightarrow \text{porphyrin})$
Zn–PM2a	$0.67 \pm 0.08$	
Zn–P3Ra	$0.32 \pm 0.03$	$0.16 \pm 0.02$
Zn–PM2b	$0.88 \pm 0.11$	
Zn–P3Rb	$0.22 \pm 0.03$	$0.10 \pm 0.03$
Zn–PM4b	$0.68 \pm 0.06$	
Zn–P5Rb	$0.25 \pm 0.02$	$0.09 \pm 0.02$

The comparison of macrocycle  $\rightarrow$  porphyrin EET efficiencies in the presence and absence of polyene show that the polyene acts as a sink for singlet excited state energy from the phenanthroline macrocycle, even though it does not quench the porphyrin.<sup>18</sup> The excited-state energy of the macrocycle is split between the porphyrin and the polyene. The higher efficiency of polyene  $\rightarrow$  porphyrin EET in [3]rotaxane Zn–P3Ra, compared with Zn–P3Rb and Zn–P5Rb, reflects the shorter polyene–porphyrin distance. The polyene  $\rightarrow$  porphyrin EET must be very rapid to compete with intersystem crossing ( $\tau \approx 0.5 \text{ ns}$ ).<sup>18</sup> Furthermore, the  $S_1$  of the polyene is a dark state with negligible oscillator strength; therefore, Förster-type energy transfer is expected to be very inefficient, implying that the observed EET occurs via a Dexter mechanism.

In conclusion, we have prepared a series of porphyrin–polyene [3]- and [5]rotaxanes that comprise multiple chromophores in a unique interlocked architecture. An analysis

of the absorption, fluorescence, and excitation spectra of the rotaxanes reveals three types of energy-transfer processes (a) from the phenanthroline macrocycle to the porphyrin, (b) from the phenanthroline macrocycle to the polyene, and (c) from the polyene to the porphyrin. This work paves the way for a template-directed synthesis of polycatenane cyclo[*n*]carbons.

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acs.orglett.6b03528](https://doi.org/10.1021/acs.orglett.6b03528).

Details of synthetic procedures and characterization data, analysis of absorption, fluorescence and excitation spectra, and crystallographic analysis (PDF)  
Crystallographic data for P3Ra (CIF)

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: [harry.anderson@chem.ox.ac.uk](mailto:harry.anderson@chem.ox.ac.uk).

### ORCID

Harry L. Anderson: [0000-0002-1801-8132](https://orcid.org/0000-0002-1801-8132)

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

We thank the EPSRC and the European Research Council (Grant No. 320969) for support, the EPSRC UK Mass Spectrometry Facility at Swansea University for mass spectra, and Diamond Light Source for an award of beamtime.

## ■ REFERENCES

- (1) (a) Diederich, F.; Rubin, Y.; Knobler, C. B.; Whetten, R. L.; Schriver, K. E.; Houk, K. N.; Li, Y. *Science* **1989**, *245*, 1088. (b) Rubin, Y.; Diederich, F. *J. Am. Chem. Soc.* **1989**, *111*, 6870. (c) Rubin, Y.; Knobler, C. B.; Diederich, F. *J. Am. Chem. Soc.* **1990**, *112*, 1607. (d) Rubin, Y.; Kahr, M.; Knobler, C. B.; Diederich, F.; Wilkins, C. L. *J. Am. Chem. Soc.* **1991**, *113*, 495. (e) McElvany, S. W.; Ross, M. M.; Goroff, N. S.; Diederich, F. *Science* **1993**, *259*, 1594. (f) Diederich, F.; Rubin, Y.; Chapman, O. L.; Goroff, N. S. *Helv. Chim. Acta* **1994**, *77*, 1441.
- (2) (a) Tobe, Y.; Fujii, T.; Matsumoto, H.; Naemura, K.; Achiba, Y.; Wakabayashi, T. *J. Am. Chem. Soc.* **1996**, *118*, 2758. (b) Tobe, Y.; Matsumoto, H.; Naemura, K.; Achiba, Y.; Wakabayashi, T. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 1800. (c) Tobe, Y.; Fujii, T.; Matsumoto, H.; Tsumuraya, K.; Noguchi, D.; Nakagawa, N.; Sonoda, M.; Naemura, K.; Achiba, Y.; Wakabayashi, T. *J. Am. Chem. Soc.* **2000**, *122*, 1762. (d) Tobe, Y.; Umeda, R.; Iwasa, N.; Sonoda, M. *Chem. - Eur. J.* **2003**, *9*, 5549.
- (3) Adamson, G. A.; Rees, C. W. *J. Chem. Soc., Perkin Trans. 1* **1996**, 1535.
- (4) von Helden, G.; Gotts, N. G.; Bowers, M. T. *Nature* **1993**, *363*, 60.
- (5) Crowley, J. D.; Goldup, S. M.; Lee, A. L.; Leigh, D. A.; McBurney, R. T. *Chem. Soc. Rev.* **2009**, *38*, 1530.
- (6) (a) Saito, S.; Takahashi, E.; Nakazono, K. *Org. Lett.* **2006**, *8*, 5133. (b) Sato, Y.; Yamasaki, R.; Saito, S. *Angew. Chem., Int. Ed.* **2009**, *48*, 504. (c) Hayashi, R.; Wakatsuki, K.; Yamasaki, R.; Mutoh, Y.; Kasama, T.; Saito, S. *Chem. Commun.* **2014**, *50*, 204. (d) Yamashita, Y.; Mutoh, Y.; Yamasaki, R.; Kasama, T.; Saito, S. *Chem. - Eur. J.* **2015**, *21*, 2139.
- (7) (a) Berná, J.; Goldup, S. M.; Lee, A.-L.; Leigh, D. A.; Symes, M. D.; Teobaldi, G.; Zerbetto, F. *Angew. Chem., Int. Ed.* **2008**, *47*, 4392. (b) Goldup, S. M.; Leigh, D. A.; Long, T.; McGonigal, P. R.; Symes, M. D.; Wu, J. *J. Am. Chem. Soc.* **2009**, *131*, 15924.
- (8) Langton, M. J.; Matichak, J. D.; Thompson, A. L.; Anderson, H. L. *Chem. Sci.* **2011**, *2*, 1897.
- (9) Movsisyan, L. D.; Kondratuk, D. V.; Franz, M.; Thompson, A. L.; Tykwinski, R. R.; Anderson, H. L. *Org. Lett.* **2012**, *14*, 3424.
- (10) (a) Weisbach, N.; Baranova, Z.; Gauthier, S.; Reibenspies, J. H.; Gladysz, J. A. *Chem. Commun.* **2012**, *48*, 7562. (b) Baranová, Z.; Amini, H.; Bhuvanesh, N.; Gladysz, J. A. *Organometallics* **2014**, *33*, 6746.
- (11) Movsisyan, L. D.; Franz, M.; Hampel, F.; Thompson, A. L.; Tykwinski, R. R.; Anderson, H. L. *J. Am. Chem. Soc.* **2016**, *138*, 1366.
- (12) Franz, M.; Januszewski, J. A.; Wendinger, D.; Neiss, C.; Movsisyan, L. D.; Hampel, F.; Anderson, H. L.; Görling, A.; Tykwinski, R. R. *Angew. Chem., Int. Ed.* **2015**, *54*, 6645.
- (13) Schrettl, S.; Contal, E.; Hoheisel, T.; Fritzsche, M.; Balog, S.; Szilluweit, R.; Frauenrath, H. *Chem. Sci.* **2015**, *6*, 564.
- (14) (a) Roche, C.; Sour, A.; Niess, F.; Heitz, V.; Sauvage, J. P. *Eur. J. Org. Chem.* **2009**, *2009*, 2795. (b) Roche, C.; Sauvage, J.-P.; Sour, A.; Strutt, N. L. *New J. Chem.* **2011**, *35*, 2820.
- (15) (a) Megiatto, J. D.; Schuster, D. I. *Org. Lett.* **2011**, *13*, 1808. (b) Megiatto, J. D.; Schuster, D. I.; de Miguel, G.; Wolfrum, S.; Guldi, D. M. *Chem. Mater.* **2012**, *24*, 2472.
- (16) Chalifoux, W.; Tykwinski, R. R. *Nat. Chem.* **2010**, *2*, 967.
- (17) Single-crystal X-ray diffraction data for Zn-P3Ra were collected with synchrotron radiation using I19 (EH1) at Diamond Light Source ( $\lambda = 0.6889 \text{ \AA}$ ): Nowell, H.; Barnett, S. A.; Christensen, K. E.; Teat, S. J.; Allan, D. R. *J. Synchrotron Radiat.* **2012**, *19*, 435. Cell parameters were determined and refined and raw frame data were integrated using CrysAlisPro (Agilent Technologies, 2010). The structure was solved with SuperFlip: Palatinus, L.; Chapuis, G. *J. Appl. Crystallogr.* **2007**, *40*, 786. It was then refined by full-matrix least-squares on  $F^2$  using CRYSTALS: Betteridge, P. W.; Carruthers, J. R.; Cooper, R. I.; Prout, K.; Watkin, D. J. *J. Appl. Crystallogr.* **2003**, *36*, 1487. Cooper, R. I.; Thompson, A. L.; Watkin, D. J. *J. Appl. Crystallogr.* **2010**, *43*, 1100. Parois, P.; Cooper, R. I.; Thompson, A. L. *Chem. Cent. J.* **2015**, *9*, 30. The structure contains large solvent-accessible voids comprising weak, diffuse electron density which were treated using PLATON/SQUEEZE: Spek, A. *J. Appl. Crystallogr.* **2003**, *36*, 7. van der Sluis, P.; Spek, A. L. *Acta Crystallogr., Sect. A: Found. Crystallogr.* **1990**, *46*, 194. Single Crystal Data:  $C_{301}H_{348}N_8O_{17}Zn$ ,  $M_r = 4415.49$ , monoclinic,  $Pc$ ;  $a = 31.7209(5) \text{ \AA}$ ,  $b = 19.3373(2) \text{ \AA}$ ,  $c = 50.4748(7) \text{ \AA}$ ,  $\beta = 96.0619(14)^\circ$ ,  $V = 30787.9(7) \text{ \AA}^3$ , data/restraints/parameters: 52640/18794/5933,  $R_{int} = 6.99\%$ , final  $R_1 = 12.36\%$ ,  $wR_2 = 31.63\%$  ( $I > 2\sigma(I)$ ),  $\Delta\rho_{min,max} = -0.78, +0.88 \text{ e \AA}^{-3}$ . Crystallographic data (excluding structure factors) have been deposited with the Cambridge Crystallographic Data Centre (CCDC 1519120); copies of these data can be obtained free of charge via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).
- (18) Movsisyan, L. D.; Peeks, M. D.; Greetham, G. M.; Towrie, M.; Thompson, A. L.; Parker, A. W.; Anderson, H. L. *J. Am. Chem. Soc.* **2014**, *136*, 17996.
- (19) Lucotti, A.; Tommasini, M.; Fazzi, D.; Del Zoppo, M.; Chalifoux, W. A.; Ferguson, M. J.; Zerbi, G.; Tykwinski, R. R. *J. Am. Chem. Soc.* **2009**, *131*, 4239.
- (20) (a) Szafert, S.; Gladysz, J. A. *Chem. Rev.* **2003**, *103*, 4175. (b) Szafert, S.; Gladysz, J. A. *Chem. Rev.* **2006**, *106*, PR1.